

Supporting Information

Boron subphthalocyanines bearing non-steroidal anti-inflammatory drug diclofenac: synthesis and photochemical properties

Tatiana V. Dubinina^{a@}, Ivan D. Burtsev^b, Alina S. Agranat,^a Anton E. Egorov,^b Alexey A. Kostyukov,^b Vladimir A. Kuzmin,^b and Elena R. Milaeva^a

^a *Department of Chemistry, Lomonosov Moscow State University, 119991 Moscow, Russian Federation.*

^b *Emanuel Institute of Biochemical Physics, Russian Academy of Sciences, 119334, Moscow, Russian Federation.*

[@] Corresponding author E-mail: dubinina.t.vid@gmail.com

Table S1. High-resolution mass spectrometry MALDI TOF/TOF data.

Figure S1. MALDI TOF mass spectrum (positive ion mode) of boron subphthalocyanine **2b**, isotopic pattern (inset A) and simulated MS pattern (inset B).

Figure S2. ^1H NMR spectrum of subphthalocyanine **2b** in CDCl_3 .

Figure S3. ^{11}B NMR spectra of subphthalocyanines **2a,b** in CDCl_3 .

Figure S4. UV-Vis spectra of subphthalocyanines **2a,b** (dashed lines) and **1a,b** (solid lines) in C_6H_6 .

Table S1. High-resolution mass spectrometry MALDI TOF/TOF data.

Compound (Molecular formula)	Mass found (ion)	Monoisotopic mass calculated
2a ($C_{74}H_{46}BCl_2N_7O_2$)	1145.3171 ($[M]^-$)	1145.3183
2b ($C_{74}H_{46}BCl_2N_7O_8$)	1241.2846 ($[M]^-$)	1241.2878




Figure S1. MALDI TOF mass spectrum (positive ion mode) of boron subphthalocyanine **2b**, isotopic pattern (inset A) and simulated MS pattern (inset B).




Figure S2. ^1H NMR spectrum of subphthalocyanine **2b** in CDCl_3 .





Figure S3. ^{11}B NMR spectra of subphthalocyanines **2a,b** in CDCl_3 .




Figure S4. UV-Vis spectra of subphthalocyanines **2a,b** (dashed lines) and **1a,b** (solid lines) in C_6H_6 .